

Opsovit® ZI

Iron, Vitamin B-Complex & Zinc

Description

Opsovit® ZI is a combination of Iron (III) Hydroxide Polymaltose Complex, Thiamine, Riboflavin, Pyridoxine, Nicotinamide & Zinc, that is used mainly in Iron or Vitamin B-Complex or Zinc deficiency states.

Composition

Opsovit® ZI Syrup: Each 5 ml syrup contains Elemental Iron 50 mg as Iron (III) Hydroxide Polymaltose Complex INN, Thiamine Hydrochloride BP 5 mg, Riboflavin 2 mg as Riboflavin Sodium Phosphate BP, Pyridoxine Hydrochloride BP 2 mg, Nicotinamide BP 20 mg & Elemental Zinc 10 mg as Zinc Sulphate Monohydrate USP.

Indications

Opsovit® ZI is indicated for the treatment and prevention of Iron or Vitamin B-Complex or Zinc deficiencies.

Dosage and administration

Adults: 5 to 10 ml (1 to 2 teaspoonful) 3 times daily.

Children: 5 ml (1 teaspoonful) 3 times daily.

Infant: 5 ml (1 teaspoonful) daily.

This may be changed as recommended by the physician.

Contraindications

Opsovit® ZI is contraindicated in patients with a known hypersensitivity to any of the ingredients of the product.

Side effects

Opsovit® ZI is generally well tolerated, although common side effects of oral Iron preparations like nausea, vomiting, constipation or diarrhoea may be observed in some cases.

Use in pregnancy and lactation

It is recommended in pregnancy and lactation.

Use in children

It is recommended for children.

Precautions

Caution should be taken in the conditions where there is a risk of iron overload, e.g. haemochromatosis, thalassemia, haemosiderosis or haemolytic anemia.

Drug interactions

No drug interaction has been reported yet. As the Iron remains as polymaltose complex, any type of ionic interaction with foodstuff components (tannins, oxalates or phytates) and medicaments (antacids, tetracycline) administered concomitantly is unlikely to occur.

Over dosage

In case of overdose, initially epigastric pain, diarrhoea and vomiting may occur. Metabolic acidosis, convulsions and coma may result after apparent recovery. Emergency medical attention is essential in the event of overdose. Primarily an emetic should be given and then gastric lavage and general supportive measures should be employed.

Storage

Keep out of reach of children. Store in a cool and dry place, protected from light.

Packaging

Opsovit® ZI Syrup: Each carton contains a bottle having 100 ml syrup with a plastic spoon.



Manufactured by
Opsonin Pharma Limited
Rupatali, Barishal, Bangladesh.
® Registered Trade Mark.

0566-01