

Indomet[®]

Indomethacin

Description

Indomethacin (**Indomet[®]**) has prominent anti-inflammatory effects with antipyretic and analgesic actions.

Mode of action

Indomethacin (**Indomet[®]**) is an inhibitor of prostaglandin synthetase. The analgesic effects result from both peripheral and central effects and are distinct from its anti-inflammatory property.

Pharmacokinetics

Following oral administration, indomethacin is rapidly and completely absorbed. Food delays absorption, but does not reduce overall bioavailability. Rectal administration produces earlier but lower peak plasma concentration. The overall bioavailability after rectal indomethacin administration is 80%. Indomethacin is 90-94% protein bound. Only very small amounts pass into the brain and into breast milk; some crosses the placenta. Indomethacin is extensively metabolized in the liver. The major route of elimination is in the urine. Only a small amount is excreted unchanged in the faeces.

Composition

Indomet[®] 25 mg Capsule: Each capsule contains Indomethacin BP 25 mg.

Indomet[®] SR 75 mg Capsule: Each capsule contains Indomethacin BP 75 mg.

Indomet[®] Suppository: Each suppository contains Indomethacin BP 100 mg.

Indications

Pain and moderate to severe inflammation in acute gout, rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, acute musculoskeletal disorders, dysmenorrhoea, closure of patent ductus arteriosus in premature babies.

Dosage & administration

Capsule: *Rheumatic disease:* 50-200 mg daily in divided doses, with food; Child not recommended.

Acute gout: 150-200 mg daily in divided doses.

Dysmenorrhoea: Up to 75 mg daily.

SR Capsule: 1 capsule 1-2 times daily; Child not recommended.

Suppository: 100 mg at night and in the morning if required; Child not recommended.

Contraindications

Indomethacin is contraindicated in children, pregnant and lactating women; active peptic ulcer; a history of recurrent gastrointestinal lesions; sensitivity to indomethacin, aspirin or other NSAIDs.

Side effects

Frequently gastrointestinal disturbances (including diarrhoea), headache, dizziness and dyspepsia. If headache persists even after dosage reduction,

indomethacin should be withdrawn. Also gastrointestinal ulceration and bleeding; rarely drowsiness, confusion, insomnia, convulsions, psychiatric disturbances, depression, syncope, hypertension, hyperglycemia and blurred vision. Suppository may cause rectal irritation and occasional bleeding.

Use in pregnancy & lactation

It is not known whether indomethacin is safe to use in pregnant and lactating women. So it should not be given to such groups.

Precautions

Indomethacin should be used with care in patients with epilepsy, parkinsonism, psychiatric disturbances. Rectal administration should be avoided in proctitis and haemorrhoids.

Drug interactions

Aspirin may decrease, and diflunisal and probenecid may increase plasma levels of indomethacin. The antihypertensive effect of beta-blockers and the diuretic and antihypertensive effects of thiazides and furosemide may be reduced by indomethacin. It reduces the tubular secretion and therefore potentiates the toxicity of methotrexate. There is an increased risk of gastrointestinal bleeding when corticosteroids are administered concomitantly with NSAIDs.

Overdosage

Nausea, vomiting, intense headache, dizziness, mental confusion, disorientation or lethargy may be observed following overdosage of indomethacin. There have been reports of paresthesias, numbness and convulsions. Treatment is symptomatic and supportive. The stomach should be emptied as quickly as possible if the ingestion is recent. If vomiting has not occurred spontaneously, the patient should be induced to vomit with syrup of ipecac. If the patient is unable to vomit, gastric lavage should be performed.

Storage

Keep out of reach of children. Store in a dry place, below 25°C temperature and protected from light.

Packaging

Indomet® 25 mg Capsule: Each carton contains 20X10 capsules in blister pack.

Indomet® SR 75 mg Capsule: Each carton contains 10X5 capsules in blister pack.

Indomet® Suppository: Each carton contains 5X5 suppositories in blister pack.



Manufactured by
Opsonin Pharma Limited
Rupatali, Barishal, Bangladesh.
® Registered Trade Mark.

1059.01