Tafost® MONO

Sterile Eye Drops

Tafluprost 0.0015%

(Preservative-free Single Vial)

Description: Tafluprost is a selective FP prostanoid receptor agonist which reduces intraocular pressure.

Mode of action: Tafluprost is a prostaglandin analog, which lowers intraocular pressure primarily by increasing uveoscleral outflow and to a lesser extent, enhancing aqueous humor drainage through the trabecular outflow pathway. In addition, Tafluprost relaxes the smooth muscle of the ciliary arteries by inhibiting Ca2+ influx into the cells. Thus, Tafluprost increases ocular blood flow and provides ocular neuroprotective action.

Composition: Tafost® Mono Preservative-free Sterile Eye Drops: Each ml contains Tafluprost INN 0.015 mg.

Indications: Indicated for reducing elevated intraocular pressure in patients with open-angle glaucoma or ocular hypertension.

Dosage & administration: Instill one drop in the affected eye(s) once daily in the evening.

Administration guidelines:

Open the sachet pack & take out the vial strip

1

Pull off one single-dose vial from the strip

2

Hold the vial upright

3

Tap the top until all liquid goes to the bottom

4

Twist off the cap without touching the tip



5

Tilt your head back and pull your lower eyelid down

6

Squeeze one drop into the lower eyelid without touching the eye

7

Close your eye and press the inner corner for one minute

8

Contraindications: Contraindicated in patients who are hypersensitive to any of the components of this preparation.

Side effects: Since Tafluprost is PRESERVATIVE-FREE, it avoids the potential risks caused by the preservatives.

Use in pregnancy & lactation: *Use in pregnancy:* Pregnancy category C.
Use in lactation: There is no well controlled study in lactating women.

Precautions: To avoid contamination or eye injury, do not allow the tip of the single vial to touch your eye or any other surface. If you wear contact lenses, remove them before instilling the ophthalmic solution and wait at least 15 minutes before reinserting them. If you are using more than one topical ophthalmic medication, allow an interval of at least 5 minutes between applications.

Drug interactions: Proper data not found.

Over dosage: Specific studies are not available.

Storage: Keep out of reach of children. Store in a dry, well ventilated place at temperature up to 2 - 8°C and protected from extraneous odors, contamination & intense light.

Packaging:
Tafost® Mono Preservative-free Sterile Eye Drops: Each carton contains 2X15 vials in Alu-Alu sachet pack. Each single vial contains 0.2 ml of sterile eye drops.

Opsonin Pharma
Ideas for healthcare

Manufactured by
Opsonin Pharma Limited
Rupatali, Barishal, Bangladesh.
® Registered Trade Mark.

টাফোস্ট® মনো

জীবাণুমুক্ত চোখের ড্রপস্‌

টাফ্লুপ্রোস্ট ০.০০১৫%

(প্রিজারভেটিভ-মুক্ত সিঙ্গেল ভায়াল)

বিবরণ: টাফ্লুপ্রোস্ট একটি নির্বাচিত এফপি প্রোস্টানয়েড রিসেপ্টর অ্যাগোনিস্ট, যা ইন্ট্রাঅকুলার প্রেসার কমায়।

কার্যপদ্ধতি: টাফ্লুপ্রোস্ট একটি প্রোস্টাগ্লান্ডিন অ্যানালগ যা প্রধানত ইউভিওস্ক্লেরাল বহিঃপ্রবাহ বৃদ্ধি করে এবং অপেক্ষাকৃত কম মাত্রায় ট্রাবেকুলার আউটফ্লোর মাধ্যমে অ্যাকুয়াস হিউমারের নিষ্কাশন করে ইন্ট্রাঅকুলার প্রেসার কমায়। এছাড়া, এটি কোষে Ca^{2+} প্রবেশ প্রতিরোধের মাধ্যমে সিলিয়ারি আর্টারির মসৃণ পেশীকে শিথিল করে। ফলে চোখে রক্তপ্রবাহ বৃদ্ধি পায় এবং নিউরোপ্রোটেক্টিভ অ্যাকশন প্রদান করে।

উপাদান: টাফোস্ট® মনো প্রিজারভেটিভ-ফ্রি জীবাণুমুক্ত চোখের ড্রপস: প্রতি মিলিতে আছে টাফ্লুপ্রোস্ট আইএনএন ০.০১৫ মিগ্রা।

নির্দেশনা: ওপেন-অ্যাঙ্গেল গ্লুকোমা অথবা অকুলার হাইপারটেনশনে আক্রান্ত রোগীদের চোখের অভ্যন্তরীণ উচ্চ চাপ কমানোর জন্য নির্দেশিত।

মাত্রা ও প্রয়োগ: আক্রান্ত চোখে প্রতি সন্ধ্যায় এক ফোঁটা করে প্রয়োগ করুন।

ব্যবহারবিধি:

স্যাশে প্যাকটি খুলুন এবং ভায়ালের স্ট্রিপটি বের করে নিন।

১

স্ট্রিপ থেকে একটি একক-ডোজের ভায়াল আলাদা করে নিন।

২

ভায়ালটি সোজাভাবে ধরে রাখুন।



৩

উপরের অংশে হালকা টোকা দিন, যতক্ষণ না সব তরল নিচের অংশে চলে আসে।

৪

টিপে স্পর্শ না করে ঢাকনাটি ঘুরিয়ে খুলে ফেলুন।



৫

আপনার মাথা পেছনের দিকে হেলান এবং নিচের চোখের পাতা নিচে টেনে ধরুন।

৬

চোখে স্পর্শ না করে নিচের চোখের পাতার ভেতরে এক ফোঁটা ড্রপ দিন।

৭

চোখ বন্ধ করুন এবং এক মিনিটের জন্য চোখের ভেতরের কোণায় চাপ দিন।

৮

বিরুদ্ধ ব্যবহার (যেসব ক্ষেত্রে ব্যবহার করা যাবে না): যেসব রোগীর এই প্রস্তুতির কোনো উপাদানের প্রতি অতিসংবেদনশীলতা রয়েছে, তাদের ক্ষেত্রে এটি ব্যবহার করা যাবে না।

পার্শ্বপ্রতিক্রিয়া: যেহেতু টাফ্লুপ্রোস্ট প্রিজারভেটিভ-মুক্ত, তাই এতে প্রিজারভেটিভজনিত সম্ভাব্য ঝুঁকি এড়ানো যায়।

গর্ভাবস্থা ও স্তন্যদানকালে ব্যবহার: গর্ভাবস্থায় ব্যবহার: প্রেগন্যান্সি ক্যাটাগরি-সি।
স্তন্যদানকালে ব্যবহার: স্তন্যদানকারী নারীদের ক্ষেত্রে এ বিষয়ে কোনো সু-নিয়ন্ত্রিত গবেষণা নেই।

সতর্কতা: দূষণ বা চোখে আঘাত এড়াতে সিঙ্গেল ভায়ালের মুখ যেন চোখ বা অন্য কোনো কিছুর সংস্পর্শে না আসে। কন্টাক্ট লেন্স ব্যবহার করলে ওষুধ প্রয়োগের আগে লেন্স খুলে ফেলুন এবং ড্রপ প্রয়োগের কমপক্ষে ১৫ মিনিট পরে পুনরায় লেন্স পরুন। একাধিক টপিক্যাল চোখের ড্রপস ব্যবহার করলে প্রতিটি ড্রপ প্রয়োগের মধ্যে অন্তত ৫ মিনিট বিরতি রাখুন।

অন্য ওষুধের সাথে প্রতিক্রিয়া: পর্যাপ্ত তথ্য পাওয়া যায়নি।

মাত্রাধিক্য: নির্দিষ্ট গবেষণা নেই।

সংরক্ষণ: শিশুদের নাগালের বাইরে রাখুন। শুষ্ক ও পর্যাপ্ত বায়ু চলাচলযুক্ত স্থানে ২°-৮° সে. পর্যন্ত তাপমাত্রায় সংরক্ষণ করুন এবং বাহ্যিক গন্ধ, দূষণ ও তীব্র আলো থেকে দূরে রাখুন।

উপস্থাপনা:
টাফোস্ট® মনো প্রিজারভেটিভ-ফ্রি জীবাণুমুক্ত চোখের ড্রপস: প্রতিটি কার্টনে রয়েছে ২×১৫ টি ভায়াল অ্যালু-অ্যালু স্যাশে প্যাকেটে। প্রতিটি সিঙ্গেল ভায়ালে আছে ০.২ মিলি জীবাণুমুক্ত চোখের ড্রপস।

Opsonin
Opsonin Pharma
Ideas for healthcare

Manufactured by
Opsonin Pharma Limited
Rupatali, Barishal, Bangladesh.
® Registered Trade Mark.

000-01